arXiv:1403.4959v1 [math.OA] 19 Mar 2014

PURELY INFINITE C^*-ALGEBRAS ASSOCIATED TO ÉTALE GROUPOIDS

JONATHAN BROWN, LISA ORLOFF CLARK, AND ADAM SIERAKOWSKI

Abstract. Let G be a Hausdorff, étale groupoid that is minimal and topologically principal. We show that $C_r^*(G)$ is purely infinite simple if and only if all the nonzero positive elements of $C_0(G^{(0)})$ are infinite in $C_r^*(G)$. If G is a Hausdorff, ample groupoid, then we show that $C_r^*(G)$ is purely infinite simple if and only if every nonzero projection in $C_0(G^{(0)})$ is infinite in $C_r^*(G)$. We then show how this result applies to k-graph C^*-algebras. Finally, we investigate strongly purely infinite groupoid C^*-algebras.

1. Introduction

Purely infinite simple C^*-algebras were introduced by Cuntz in [7] where he showed that the K_0 group of such algebras can be computed within the algebra itself without resorting to the usual direct limit construction. The K-theory groups of C^*-algebras have long been known to be computable invariants and Cuntz's result shows that this computation is easier when the C^*-algebra is purely infinite simple. Elliott initiated a program to find a suitably large class of C^*-algebras on which the K-theory groups provide a complete isomorphism invariant (see [8]). This program has achieved remarkable success, most notably in a theorem of Kirchberg and Phillips [11, 20] which states that every *Kirchberg algebra* satisfying the Universal Coefficient Theorem (UCT) is classified by the isomorphism classes of its ordered K-theory groups. A Kirchberg algebra is a separable, nuclear, purely infinite simple C^*-algebra.

The allure of classification via the Kirchberg-Phillips theorem has lead many authors to study when various constructions of C^*-algebras yield purely infinite simple algebras. Kumjian, Pask and Raeburn show that a graph C^*-algebra of a cofinal graph is purely infinite simple if and only if every vertex can be reached from a loop with an entrance [17, Theorem 3.9]. Carlsen and Thomsen show that if the C^*-algebra constructed from a locally injective surjection θ on a compact metric space of finite covering dimension is simple, then it is purely infinite simple if and only if θ is not a homeomorphism [5, Corollary 6.6]. Rørdam and Sierakowski [24] show that if a countable exact group H acts by an essentially free action on the Cantor set X and the type semigroup of clopen subsets of X is almost unperforated, then $C_0(X) \rtimes_r H$ is purely infinite if and only if every clopen set E in X is paradoxical. The constructions in each of the above examples are special cases of groupoid C^*-algebras.

Date: March 21, 2014.

2010 *Mathematics Subject Classification.* 46L05 (Primary); 22A22 (Secondary).

Key words and phrases. Groupoid; groupoid C^*-algebra; purely infinite C^*-algebra; strongly purely infinite; topologically principal; k-graph.

This research was supported by the Australian Research Council.

In this paper we investigate purely infinite C^*-algebras associated to Hausdorff, étale groupoids. In sections 3 through 5 we restrict our attention to simple groupoid C^*-algebras. Characterising simplicity of groupoid C^*-algebras is known and we readily make use of the following theorem from [4, Theorem 5.1]:

Theorem 1.1 (Brown-Clark-Farthing-Sims)**.** *Let G be a second-countable, locally compact, Hausdorff and étale groupoid. Then $C^*(G)$ is simple if and only if all of the following conditions are satisfied.*

(1) $C^*(G) = C_r^*(G)$*;*
(2) G *is topologically principal;*
(3) G *is minimal.*

However, necessary and sufficient conditions on the groupoid for the associated algebra to be *purely infinite* simple are not known. Anantharaman-Delaroche showed that 'locally contracting' is a sufficient condition on the groupoid in [1, Proposition 2.4] but whether locally contracting is necessary remains an open question. Part of the difficulty in characterizing those groupoids that give rise to purely infinite simple C^*-algebras is relating arbitrary projections in the groupoid C^*-algebra to the groupoid itself.

In Section 3 we look at a necessary and sufficient conditions for ensuring pure infiniteness of groupoid C^*-algebras. We show that for a Hausdorff, étale, topologically principal, and minimal groupoid G the C^*-algebra $C_r^*(G)$ is purely infinite simple if and only if it all the nonzero positive elements of $C_0(G^{(0)})$ are infinite in $C_r^*(G)$ and (see Theorem 3.3). In Section 4 we specialize to Hausdorff, *ample* groupoids. This is an important class of examples because every Kirchberg algebra in UCT is Morita equivalent to an algebra associated to an Hausdorff, ample groupoid (see [26]). We show in Theorem 4.1 for a Hausdorff, ample groupoid G, that is also topologically principal and minimal, the C^*-algebra $C_r^*(G)$ is purely infinite if and only if every nonzero projection in $C_0(G^{(0)})$ is infinite in $C_r^*(G)$. Theorem 4.1 is a generalisation of [10] about partial actions. In Section 5 we demonstrate how Theorem 4.1 applies to k-graph C^*-algebras.

In Section 6 we turn our attention to the non-simple case. In [13], Kirchberg and Rørdam introduce three separate notions of purely infinite C^*-algebras: weakly purely infinite, purely infinite and strongly purely infinite. Of these notions, the last one appears to be the most useful in the classification theory of non-simple C^*-algebras. Indeed, Kirchberg showed in [12] that two separable, nuclear, strongly purely infinite C^*-algebras with the same primitive ideal space X are isomorphic if and only if they are KK_X-equivalent. We provide a characterization of when groupoid C^*-algebras are strongly purely infinite in Proposition 6.3.

2. Preliminaries

2.1. **Groupoids.** A *groupoid* G is a small category in which every morphism is invertible. The set of objects in G is identified with the set of identity morphisms and both are denoted by $G^{(0)}$. We call $G^{(0)}$ the *unit space* of G. Each morphism γ in the category has a range and source denoted $r(\gamma)$ and $s(\gamma)$ respectively and thus r and s define maps $G \to G^{(0)}$.

A *topological groupoid* is a groupoid with a topology in which composition is continuous and inversion is a homeomorphism. An *open bisection* in a topological groupoid G is an open set B such that both r and s restricted to B are homeomorphisms; in particular

these restrictions are injective. An *étale groupoid* is a topological groupoid where s is a local homeomorphism. If a groupoid G is Hausdorff and étale, then the unit space $G^{(0)}$ is open and closed in G. If G is a locally compact, Hausdorff groupoid, then G is étale if and only if there is a basis for the topology on G consisting of open bisections with compact closure. A topological groupoid is called *ample* if it has a basis of compact open bisections. If G is locally compact, Hausdorff and étale groupoid, then we note that G is ample if and only if $G^{(0)}$ is totally disconnected (see [9, Proposition 4.1]).

For a subsets $L, K \subseteq G$, denote $LK = \{\gamma : \gamma = \xi\zeta \quad \text{with } \xi \in L, \zeta \in K, s(\xi) = r(\zeta)\}$. With a slight abuse of notation, for $u \in G^{(0)}$, we write uG and Gu for $\{u\}G$ and $G\{u\}$ respectively and denote by uGu the set

$$\{\gamma \in G : r(\gamma) = s(\gamma) = u\}.$$

A topological groupoid G is *topologically principal* if the set $\{u \in G^{(0)} : uGu = \{u\}\}$ is dense in $G^{(0)}$, and *minimal* if $G \cdot u := \{r(\gamma) : s(\gamma) = u\}$ is dense in $G^{(0)}$ for all $u \in G^{(0)}$. Recall, for a second countable, locally compact, Hausdorff, étale groupoid G the algebra $C^*(G)$ is simple if and only if G is minimal, topologically principal, and $C^*(G) = C_r^*(G)$.

2.2. **Groupoid C^*-algebras.** Let G be locally compact, Hausdorff étale groupoid and let $C_c(G)$ denote the set of compactly supported continuous functions from G to $\mathbb{C}$. Since every element γ of G has a neighbourhood B_γ such that $r|_{B_\gamma}$ is injective, the set $r^{-1}(u)$ is discrete for every $u \in G^{(0)}$. Thus if $f \in C_c(G)$ then $\text{supp}(f) \cap r^{-1}(u)$ is finite for all $u \in G^{(0)}$. With this, we are able to define a convolution and involution on $C_c(G)$ such that for $f, g \in C_c(G)$,

$$f * g(\gamma) := \sum_{r(\eta)=r(\gamma)} f(\eta)g(\eta^{-1}\gamma) \quad \text{and} \quad f^*(\gamma) := \overline{f(\gamma^{-1})}.$$

Under these operations, $C_c(G)$ is a $*$-algebra. Next define for $f \in C_c(G)$,

$$\|f\|_I := \sup_{u \in G^{(0)}} \{\max\{\sum_{\gamma \in Gu} |f(\gamma)|, \sum_{\gamma \in uG} |f(\gamma)|\}\} \quad \text{and}$$
$$\|f\| := \sup\{\|\pi(f)\| : \pi \text{ is a } \|\cdot\|_I\text{-decreasing representation}\}.$$

Then $C^*(G)$ is the completion of $C_c(G)$ in the $\|\cdot\|$-norm.

Given a unit $u \in G^{(0)}$, the regular representation π_u of $C_c(G)$ on $\ell^2(Gu)$ associated to u is characterized by

$$\pi_u(f)\delta_\gamma = \sum_{s(\eta)=r(\gamma)} f(\eta)\delta_{\eta\gamma}.$$

The reduced C^*-norm on $C_c(G)$ is $\|f\|_r = \sup\{\|\pi_u(f)\| : u \in G^{(0)}\}$ and $C_r^*(G)$ is the completion of $C_c(G)$ in the $\|\cdot\|_r$-norm. Our attention will be focused on the reduced C^*-algebra and situations where the reduced and full algebras coincide. Also, we will often be assuming G is second-countable, implying $C_r^*(G)$ is separable [21, Page 59].

Below we recall a few standard results (we heavily use) and their proofs to familiarise the reader with locally compact, Hausdorff étale groupoids.

Lemma 2.1 (cf. [21])**.** *Let G be a locally compact, Hausdorff and étale groupoid. Then*

(1) *The extension map from $C_c(G^{(0)})$ into $C_c(G)$ (where a function is defined to be zero on $G - G^{(0)}$) extends to an embedding of $C_0(G^{(0)})$ into $C_r^*(G)$.*

(2) *The restriction map* $E_0 : C_c(G) \to C_c(G^{(0)})$ *extends to a conditional expectation*[1] $E : C_r^*(G) \to C_0(G^{(0)})$.
(3) *The map* E *from item (2) is faithful. That is,* $E(b^*b) = 0$ *implies* $b = 0$ *for* $b \in C_r^*(G)$.
(4) *For every closed invariant set* $D \subseteq G^{(0)}$ *we have the following commuting diagram:*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C_r^*(G|_U) & \xrightarrow{\iota_r} & C_r^*(G) & \xrightarrow{\rho_r} & C_r^*(G|_D) & \longrightarrow & 0\ , \\ & & \downarrow{\scriptstyle E_U} & & \downarrow{\scriptstyle E} & & \downarrow{\scriptstyle E_D} & & \\ 0 & \longrightarrow & C_0(U) & \xrightarrow{\iota_0} & C_0(G^{(0)}) & \xrightarrow{\rho_0} & C_0(D) & \longrightarrow & 0 \end{array}$$

where $U = G^{(0)} - D$, ι_r *and* ρ_r *are determined on continuous functions by extension and restriction respectively. Moreover,* $\mathrm{image}(\iota_r) \subseteq \ker \rho_r$.
(5) *The subalgebra* $C_c(G^{(0)})$ *contains an approximate unit for* $C_r^*(G)$.

Proof. (1) Since G is Hausdorff and étale, $G^{(0)}$ is open and closed in G. Thus, the map $C_c(G^{(0)})$ into $C_c(G)$ is well defined. For $f, g \in C_c(G^{(0)})$, a quick computation gives

$$f * g(\gamma) = \begin{cases} f(\gamma)g(\gamma), & \text{if} \quad \gamma \in G^{(0)}; \\ 0, & \text{otherwise,} \end{cases}$$

so the map from $C_c(G^{(0)})$ into $C_c(G)$ is a *-homomorphism. We claim the map is isometric, that is, we claim the reduced norm agrees with the infinity norm for functions in $C_c(G^{(0)})$. By evaluating at point masses in $\ell^2(Gu)$, one can show that $\|f\|_\infty \leq \|f\|_r$ for $f \in C_c(G)$. The reverse inequality can be verified for $f \in C_c(G^{(0)})$ and the claim follows. Thus the *-homomorphism from $C_c(G^{(0)})$ into $C_c(G) \subseteq C_r^*(G)$ extends by continuity to an isometric (hence injective) *-homomorphism from $C_0(G^{(0)})$ into $C_r^*(G)$.

(2) Once again using that G is Hausdorff and ètale, we have that $G^{(0)}$ is open and closed in G and hence E_0 is well defined. One may easily verify that E_0 is (a) positive (b) linear (c) idempotent, and (d) of norm one. Therefore E_0 extends by continuity to a map $E : C_r^*(G) \to C_0(G^{(0)})$ with the same properties (a)–(d). By [3, II.6.10.1] we conclude that E is a conditional expectation.

(3) Let $b \in C_r^*(G)$ such that $E(b^*b) = 0$. We need to show that $b = 0$. Let $V_\gamma : \mathbb{C} \to \ell^2(G_{s(\gamma)})$ be given by $c \mapsto c\delta_\gamma$. Then $V_\gamma^*\omega = \omega(\gamma)$. Since $\|b\|_r = \sup_{u \in G^{(0)}} \|\pi_u(b)\|$ and

$$\|\pi_u(b)\delta_\gamma\|^2 = \langle \pi_u(b)\delta_\gamma, \pi_u(b)\delta_\gamma \rangle = \langle \pi_u(b^*b)\delta_\gamma, \delta_\gamma \rangle = V_\gamma^*\pi_u(b^*b)V_\gamma 1,$$

it suffices to show that $V_\gamma^*\pi_u(b^*b)V_\gamma = 0$ for all $u \in G^{(0)}$ and $\gamma \in G$.

For $f \in C_c(G)$, $u \in G^{(0)}$, and $c \in \mathbb{C}$, we have

$$V_u^*\pi_u(f)V_u c = V_u^*\pi_u(f)c\delta_u = V_u^*\Big(\sum_{s(\eta)=u} f(\eta)c\delta_\eta\Big) = f(u)c = E(f)(u)c. \tag{2.1}$$

Thus by the continuity of E, for all $a \in C_r^*(G)$, $E(a)(u) = V_u^*\pi_u(a)V_u$ as operators on $\mathbb{C}$.

For every open bisection B and $\gamma \in B$, pick a function $\phi_{\gamma,B} \in C_c(G)$ such that $\phi_{\gamma,B}(\gamma) = 1$, $\mathrm{supp}(\phi_{\gamma,B}) \subseteq B$, and $0 \leq \phi_{\gamma,B} \leq 1$. Now if $f \in C_c(G)$ and B is an open bisection with

[1]Recall: A *conditional expectation* $E\colon A \to B$ is a contractive, linear, completely positive map such that for every $b \in B, a \in A$ we have $E(b) = b$, $E(ba) = bE(a)$ and $E(ab) = E(a)b$, see [3, II.6.10.1].

$\gamma \in B$, then

$$(E(\phi_{\gamma,B}^* f \phi_{\gamma,B}))(u) = \sum_{r(\xi)=r(\zeta)=u} \phi_{\gamma,B}(\xi^{-1}) f(\xi^{-1}\zeta)\phi_{\gamma,B}(\zeta^{-1}),$$

which is zero unless $\xi, \zeta \in B^{-1}$. Since $r(\xi) = r(\zeta) = u$, we have that $\xi = \zeta$ is the unique element of uB^{-1}. So

$$(E(\phi_{\gamma,B}^* f \phi_{\gamma,B}))(u) = \phi_{\gamma,B}(\zeta^{-1}) f(s(\zeta))\phi_{\gamma,B}(\zeta^{-1}) \leq E(f)(s(\zeta)) \leq \|E(f)\|_\infty. \tag{2.2}$$

Now if $a \in C_r^*(G)$ then $\phi_{\gamma,B}^* a^* a \phi_{\gamma,B}$ is positive so $E(\phi_{\gamma,B}^* a^* a \phi_{\gamma,B}) \geq 0$. Therefore by the continuity of E we can apply (2.2) to obtain

$$0 \leq E(\phi_{\gamma,B}^* b^* b \phi_{\gamma,B}) \leq \|E(b^* b)\|_\infty = 0.$$

Thus $E(\phi_{\gamma,B}^* b^* b \phi_{\gamma,B}) = 0$ for all open bisections B and $\gamma \in B$.

For $\gamma \in G$ pick an open bisection B such that $\gamma \in B$. Notice for $c \in \mathbb{C}$

$$\pi_{s(\gamma)}(\phi_{\gamma,B}) V_{s(\gamma)} c = \pi_{s(\gamma)}(\phi_{\gamma,B}) c \delta_{s(\gamma)} = \sum_{s(\eta)=s(\gamma)} \phi_{\gamma,B}(\eta) c \delta_\eta = c\delta_\gamma = V_\gamma c.$$

Thus $\pi_{s(\gamma)}(\phi_{\gamma,B}) V_{s(\gamma)} = V_\gamma$ as operators. Now by equation (2.1) and the above observation we get for all $\gamma \in G$ that

$$V_\gamma^* \pi_u(b^* b) V_\gamma = V_{s(\gamma)}^* \pi_{s(\gamma)}(\phi_{\gamma,B}^* b^* b \phi_{\gamma,B}) V_{s(\gamma)}^* = E(\phi_{\gamma,B}^* b^* b \phi_{\gamma,B}) = 0$$

as desired. Therefore $b = 0$ and hence E is faithful.

(4) The diagram commutes when restricting to continuous functions with compact support. Commutativity then passes to the respective completions by continuity. Since we know $\rho_r(\iota_r(f)) = 0$ for all $f \in C_c(G|_U)$ we obtain $\text{image}(\iota_r) \subseteq \ker \rho_r$ by continuity.

(5) Let $\mathcal{C}$ be the set of compact sets in $G^{(0)}$ ordered by inclusion. For each $C \in \mathcal{C}$ pick a function e_C in $C_c(G^{(0)})$ such that $0 \leq e_c \leq 1$ and $e_C|_C \equiv 1$. Fix $f \in C_c(G)$ vanishing outside a compact set $K \subseteq G$. For C such that $s(K) \subset C$, $f * e_c = f$. It follows that (e_C) is an approximate unit for $C_r^*(G)$. □

2.3. **Purely infinite simple C^*-algebras.** Given a C^*-algebra A we denote its positive elements by A^+. If B is a subalgebra of A then $B^+ \subseteq A^+$. In particular, if $C_0(X)$ is an abelian subalgebra of A and $f \in C_0(X)$ such that $f(x) \geq 0$ for all $x \in X$, then $f \in A^+$.

For positive elements $a \in M_n(A)$ and $b \in M_m(A)$, a is *Cuntz below* b, denoted $a \precsim b$, if there exists a sequence of elements x_k in $M_{m,n}(A)$ such that $x_k^* b x_k \to a$ in norm. Notice that $\precsim$ is transitive: if $a \precsim b$ and $b \precsim c$ there exist sequences of element x_n and y_n such that $x_n^* b x_n \to a$ and $y_n^* c y_n \to b$ in norm, so $x_n^* y_n^* c y_n x_n \to a$ in norm, that is $a \precsim c$. We say A is *purely infinite* if there are no characters on A and for all $a, b \in A^+$, $a \precsim b$ if and only if $a \in \overline{AbA}$ [14, Definition 4.1]. A nonzero positive element $a \in A$ is *properly infinite* if $a \oplus a \precsim a$. By [14, Theorem 4.16] A is purely infinite if and only if every nonzero positive element in A is properly infinite.

A projection p in a C^*-algebra A is *infinite* if it is Murray-von Neumann equivalent to a proper subprojection of itself, i.e., if there exists a partial isometry s such that $s^* s = p$ but $ss^* \lneq p$. By [14, Proposition 4.7] a C^*-algebra A is purely infinite if every nonzero hereditary C^*-subalgebra in every quotient of A contains an infinite projection. For simple C^*-algebras the converse is also true, thus a simple C^*-algebra is purely infinite precisely when every hereditary subalgebra contains an infinite projection.

3. Topologically principal groupoids and positive elements of $C_0(G^{(0)})$

In this section we consider, locally compact, Hausdorff and étale groupoids. We will show that we can determine when $C_r^*(G)$ is purely infinite simple by restricting our attention to elements of $C_0(G^{(0)})$ (see Theorems 4.1 and 3.3). Before we do that, we need the following technical lemmas.

Lemma 3.1. *Let G be a locally compact, Hausdorff and étale groupoid and $E : C_r^*(G) \to C_0(G^{(0)})$ be the faithful conditional expectation extending restriction. Suppose that G is topologically principal. For every $\epsilon > 0$ and $c \in C_r^*(G)^+$, there exists $f \in C_0(G^{(0)})^+$ such that:*

(1) $\|f\| = 1$;
(2) $\|fcf - fE(c)f\| < \epsilon$;
(3) $\|fE(c)f\| > \|E(c)\| - \epsilon$.

Proof. Let $\epsilon > 0$. For $c = 0$ the result is trivial so let $c \in C_r^*(G)^+$ such that $c \neq 0$. Define

$$a := \frac{c}{\|E(c)\|}.$$

To find an appropriate f, we use the construction in the proof of [1, Proposition 2.4]; we include the details below. Find $b \in C_c(G) \cap C_r^*(G)^+$ so that $\|a - b\| < \frac{\epsilon}{2\|E(c)\|}$. Then

$$\|E(b)\| > 1 - \frac{\epsilon}{2\|E(c)\|}$$

because E is linear and $\|E(a)\| = 1$. Now, let $K := \operatorname{supp}(b - E(b))$, which is a compact subset of $G \setminus G^{(0)}$. Let

$$U := \{u \in G^{(0)} \mid E(b)(u) > 1 - \frac{\epsilon}{2\|E(c)\|}\}.$$

Since G is topologically principal, [1, Lemma 2.3] implies that there exists a nonempty open set $V \subseteq U$ such that $VKV = \emptyset$. Using regularity, fix a nonempty open set W such that $\overline{W} \subseteq V$. Using normality, select a positive (nonzero) real-valued function $f \in C_c(G^{(0)})$ such that $f|_{\overline{W}} = 1$, $\operatorname{supp}(f) \subseteq V$, and $0 \leq f(x) \leq 1$ for all $x \in G^{(0)}$. Therefore, f is positive in $C_r^*(G)$ and satisfies item (1).

To see that item (2) holds, a direct computation gives

$$fbf = fE(b)f. \tag{3.1}$$

Since $\|a - b\| < \frac{\epsilon}{2\|E(c)\|}$, $\|f\| = 1$ and E is norm decreasing we have

$$\|fE(a)f - fE(b)f\| < \frac{\epsilon}{2\|E(c)\|}. \tag{3.2}$$

Combining equations (3.1) and (3.2) we get

$$\|faf - fE(a)f\| = \|faf - fbf + fbf - fE(b)f + fE(b)f - fE(a)f\| < \frac{\epsilon}{\|E(c)\|}.$$

Thus multiplying by $\|E(c)\|$ gives $\|fcf - fE(c)f\| < \epsilon$ as needed in (2).

To see item (3) notice that since $\operatorname{supp} f \subseteq U$ we have

$$fE(b)f \geq (1 - \frac{\epsilon}{2\|E(c)\|})f^2.$$

Since $\|f\| = 1$, from the above equation and equation (3.2) we get

$$\|fE(a)f\| > \|fE(b)f\| - \frac{\epsilon}{2\|E(c)\|} \geq 1 - \frac{\epsilon}{2\|E(c)\|} - \frac{\epsilon}{2\|E(c)\|} = 1 - \frac{\epsilon}{\|E(c)\|}.$$

Multiplying by $\|E(c)\|$ we obtain $\|fE(c)f\| > \|E(c)\| - \epsilon$ as needed. □

Lemma 3.2. *Let G be a locally compact, Hausdorff and étale groupoid and $E : C_r^*(G) \to C_0(G^{(0)})$ be the faithful conditional expectation extending restriction. Suppose that G is topologically principal. For every nonzero $a \in C_r^*(G)^+$, there exists nonzero $h \in C_0(G^{(0)})^+$ such that $h \precsim a$.*

Proof. Let $a \in C_r^*(G)^+$ such that $a \neq 0$. Since E is faithful, $E(a)$ is nonzero. Applying Lemma 3.1 to $c := \frac{a}{\|E(a)\|}$ and $\epsilon = 1/4$ gives us an $f \in C_0(G^{(0)})$ such that items (1), (2) and (3) of Lemma 3.1 hold. In particular $\|fE(c)f\| > \frac{3}{4}$.

Following [14, p. 640], for each $d \in C_0(G^{(0)})^+$ we define the element

$$(d - 1/2)_+ := \phi_{1/2}(d) \in C_0(G^{(0)})^+$$

where $\phi_{1/2}(t) = \max\{t - 1/2, 0\}$ for $t \in \mathbb{R}^+$. Notice that

$$\|\phi_{1/2}(d)\| = \max\{\|d\| - 1/2, 0\},$$

for each $d \in C_0(G^{(0)})^+$.

Now let $h := (fE(c)f - 1/2)_+ \in C_0(G^{(0)})^+$. Using item (2) of Lemma 3.1 and [13, Lemma 2.2], we can find $g \in C_r^*(G)$ so that $h = g^* fcfg$. Therefore $h \precsim a$. Finally, $h \neq 0$ since

$$\|h\| = \|(fE(c)f - 1/2)_+\| \geq \|fE(c)f\| - 1/2 \geq 1/4 > 0.$$ □

We are now in a position to prove the main result of this section.

Theorem 3.3. *Let G be a locally compact, Hausdorff and étale groupoid. Suppose that G is minimal and topologically principal. Then $C_r^*(G)$ is purely infinite if and only if every nonzero positive element of $C_0(G^{(0)})$ is infinite in $C_r^*(G)$.*

Proof. The forward implication is trivial. To see the reverse, let $a \in C_r^*(G)^+$ such that $a \neq 0$. Using Lemma 3.2 we can find a nonzero $h \in C_0(G^{(0)})^+$ such that $h \precsim a$. By assumption, we know h is infinite. Since $C_r^*(G)$ is simple by [21, Proposition II.4.6], h is properly infinite by [14, Proposition 3.14]. Thus a is properly infinite by [14, Lemma 3.8], hence $C_r^*(G)$ is purely infinite. □

Recall that a Kirchberg algebra is a separable, nuclear, purely infinite simple C^*-algebra. We combine Theorem 3.3 with results from [2, 4, 21] to obtain the following characterization of groupoid Kirchberg algebras.

Corollary 3.4. *Let G be a second-countable, locally compact, Hausdorff and étale groupoid. Then $C^*(G)$ is a Kirchberg algebra if and only if G is minimal, topologically principal, measure-wise amenable and every non-zero positive element of $C_0(G^{(0)})$ is infinite in $C^*(G)$.*

Proof. Suppose $C^*(G)$ is a Kirchberg algebra. Then $C^*(G)$ is simple by definition and so $C^*(G) = C_r^*(G)$, G is minimal and G topologically principal [4, Theorem 5.1]. Since $C^*(G)$ is nuclear, $C_r^*(G)$ is also nuclear hence G is measure-wise amenable by [2, Corollary 6.2.14]. Finally, we apply Theorem 3.3 to see that every non-zero positive element of $C_0(G^{(0)})$ is infinite in $C^*(G)$.

Conversely, suppose G is minimal, topologically principal, measure-wise amenable and that every non-zero positive element of $C_0(G^{(0)})$ is infinite in $C^*(G)$. Then $C_r^*(G) = C^*(G)$ is nuclear by [2, Corollary 6.2.14], simple by [4, Theorem 5.1], separable because G is second countable [21, Remark (iii) page 59] and purely infinite by Theorem 3.3. □

4. C^*-algebras associated to ample groupoids

In this section, we will restrict our attention to ample groupoids. Although this might seem a very restrictive class of groupoids, it actually includes a lot of important examples. Again, every Kirchberg algebra in UCT is Morita equivalent to a C^*-algebra associated to a Hausdorff, ample groupoid (see [26]). The ample case is far more manageable than the general case. In particular there is a large number of projections in the associated algebra. Let G be a locally compact, Hausdorff and étale groupoid. If G is ample, then the complex *Steinberg algebra* associated to G is

$$A(G) := \operatorname{span}\{\chi_B : B \text{ is a compact open bisection}\} \subseteq C_c(G)$$

where χ_B denotes the characteristic function of B, is dense in $C_r^*(G)$ see [6, Proposition 4.2] (see also [27]). A quick computation shows that $\chi_B * \chi_D = \chi_{BD}$ and $\chi_B^* = \chi_{B^{-1}}$, so that if $B \subseteq G^{(0)}$ is compact open, then χ_B is a projection.

Theorem 4.1. *Let G be a second countable, Hausdorff, ample groupoid. Suppose that G is topologically principal, minimal and that $\mathcal{B}$ is a basis of $G^{(0)}$ consisting of compact open sets. Then $C_r^*(G)$ is purely infinite if and only if every nonzero projection p in $C_0(G^{(0)})$ with $\operatorname{supp}(p) \in \mathcal{B}$ is infinite in $C_r^*(G)$.*

Proof. The forward implication is trivial. To see the reverse, suppose every nonzero projection p of $C_0(G^{(0)})$ with $\operatorname{supp}(p) = U$ for some $U \in \mathcal{B}$ is infinite in $C_r^*(G)$. By Theorem 3.3 it suffices to show that every positive element in $C_0(G^{(0)})^+$ is infinite. Let $a \in C_r^*(G^{(0)})^+$ be a nonzero element. We show that a is properly infinite. We claim there is a nonzero projection $p \in C_0(G^{(0)})^+$ with $\operatorname{supp}(p) \subseteq U$ for some $U \in \mathcal{B}$ such that $p \precsim a$. To see this, first note that characteristic functions of the form χ_V are projections in $C_0(G^{(0)})$ for every compact open set $V \subseteq G^{(0)}$. Since $\mathcal{B}$ is a basis of compact open sets, there exists a compact open set $U_0 \in \mathcal{B}$ and a nonzero $s \in \mathbb{R}^+$ such that $\chi_{U_0}(x) \leq sa(x)$ for every $x \in G^{(0)}$. Then $p := \chi_{U_0} \leq sa$. Applying [14, Proposition 2.7] we get that $p \precsim sa$ and so $p \precsim a$ as claimed. Since p is infinite by assumption and $C_r^*(G)$ is simple, p is properly infinite by [14, Proposition 3.14]. Hence a is properly infinite by [14, Lemma 3.8]. □

In the next corollary, we show how we can use the minimality of G to strengthen our result.

Corollary 4.2. *Let G be a second countable, Hausdorff, ample groupoid. Suppose that G is topologically principal and minimal. Then $C_r^*(G)$ is purely infinite if and only if there exists a point $x \in G^{(0)}$ and a neighbourhood basis $\mathcal{D}$ at x consisting of compact open sets so that every nonzero projection q in $C_0(G^{(0)})$ with $\operatorname{supp}(q) \in \mathcal{D}$ is infinite in $C_r^*(G)$.*

Proof. Again, the forward direction is trivial. For the reverse implication, suppose there exist a point $x \in G^{(0)}$ and neighbourhood basis $\mathcal{D}$ of x consisting of compact open sets such that that every nonzero projection q in $C_0(G^{(0)})$ with $\operatorname{supp}(q) \in \mathcal{D}$ is infinite in $C_r^*(G)$. Let $\mathcal{B}$ be a basis of $G^{(0)}$ consisting of compact open sets and suppose $p := \chi_U$ is a nonzero projection in $C_0(G^{(0)})$ with $U \in \mathcal{B}$. By Theorem 4.1, it suffices to show that p

is infinite. Since G is minimal and ample, there exists a compact open bisection B such that $x \in s(B)$ and $r(B) \cap U \neq \emptyset$. By shrinking B, we may assume that $r(B) \subseteq U$. Since $s(B)$ is an compact open neighbourhood of x, there exists a $V \in \mathcal{D}$ such that $V \subseteq s(B)$. By shrinking B again, we may assume that $s(B) = V$. Thus,

$$\chi_V = \chi_B^* \chi_{r(B)} \chi_B.$$

That is, $\chi_V \precsim \chi_{r(B)}$. Hence, $\chi_{r(B)}$ is properly infinite by [14, Lemma 3.8]. Finally, since $\chi_U = \chi_{r(B)} + \chi_{U-r(B)}$, χ_U is infinite. □

5. An application: k-graph C^*-algebras

In this section, we apply Theorem 4.1 to C^*-algebras associated to k-graphs. We assume the reader is familiar with the basic definitions and constructions of k-graphs and their C^*-algebras found in [16], but we recall a few facts here. Let Λ be a k-graph. Then the associated C^*-algebra $C^*(\Lambda)$ is the universal C^*-algebra generated by a Cuntz-Krieger Λ-family $\{s_\lambda : \lambda \in \Lambda\}$. To keep things clean, we will restrict our attention to row-finite k-graphs with no sources but similar results hold in the more general setting. We think our results will be useful in this setting because necessary and sufficient conditions on Λ for $C^*(\Lambda)$ to be purely infinite simple are not known.

Following [16] we recall how $C^*(\Lambda)$ can be realised as the C^*-algebra of a second countable, Hausdorff, ample groupoid G_Λ as follows. Let Λ^∞ denote the infinite path space of Λ and $\Lambda^\infty(v)$ be the set of infinite paths with range v. Define

$$G_\Lambda := \{(x, n, y) \in \Lambda^\infty \times \mathbb{N}^k \times \Lambda^\infty : \sigma^l(x) = \sigma^m(y), n = l - m\}$$

where σ is the shift map. We view (x, n, y) as a morphism with source y and range x. Composition is given by $(x, n, y)(y, m, w) = (x, n+m, w)$. The unit space $G_\Lambda^{(0)}$ is identified with Λ^∞. For $\lambda, \mu \in \Lambda$ with $s(\lambda) = s(\mu)$ we define

$$Z(\lambda, \mu) := \{(\lambda z, d(\lambda) - d(\mu), \mu z) : z \in \Lambda^\infty(s(\lambda))\}.$$

The (countable) collection of all such $Z(\lambda, \mu)$ generate a topology under which G_Λ is a second countable, Hausdorff, ample groupoid by [16, Proposition 2.8]. Further, the relative topology on the unit space Λ^∞ has a basis of compact cylinder sets

$$Z(\lambda) := \{\lambda x \in \Lambda^\infty : x \in \Lambda^\infty(s(\lambda))\}$$

by identifying $Z(\lambda, \lambda)$ and $Z(\lambda)$ from [16, Lemma 2.6 and Proposition 2.8]. Note that G_Λ is amenable by [16, Theorem 5.5] and hence $C_r^*(G_\Lambda) = C^*(G_\Lambda)$. It was shown in [16] that

$$C^*(\Lambda) \cong C^*(G_\Lambda).$$

More specifically, by [16, Corollary 3.5(i)], there is a (unique) isomorphism $\phi : C^*(\Lambda) \to C^*(G_\Lambda)$ such that $\phi(s_\lambda) = \chi_{Z(\lambda, s(\lambda))}$. Note that

$$\phi(s_\mu s_\mu^*) = \chi_{Z(\mu, s(\mu))} \chi_{Z(\mu, s(\mu))}^* = \chi_{Z(\mu, s(\mu))} \chi_{Z(s(\mu), \mu)} = \chi_{Z(\mu, \mu)} = \chi_{Z(\mu)}.$$

With all of this theory in place, along with the simplicity results of [23] and [4], the following is an immediate corollary of Theorem 4.1 and Corollary 4.2.

Corollary 5.1. *Suppose Λ is a row-finite k-graph with no sources such that Λ is aperiodic and cofinal in the sense of* [23]. *Then*

1. *For $\mu \in \Lambda$, $s_\mu s_\mu^*$ is infinite if and only if $s_{s(\mu)}$ is.*
2. *$C^*(\Lambda)$ is purely infinite simple if and only if s_v is infinite for every $v \in \Lambda^0$.*

(3) $C^*(\Lambda)$ *is purely infinite simple if and only if there exists* $x \in \Lambda^\infty$ *such that* s_v *is infinite for every vertex* v *on* x.

Proof. For (1), we use a trick used in [25, Lemma 8.13]. Recall that infiniteness is preserved under von Neumann equivalence, hence $s_\mu s_\mu^*$ is infinite if and only if $s_\mu^* s_\mu = s_{s(\mu)}$ is infinite.

For (2), we apply Theorem 4.1 to the second countable, Hausdorff, ample groupoid G_Λ; first we check the remaining hypotheses of Theorem 4.1. Since Λ is cofinal and aperiodic, $C^*(G_\Lambda) \cong C^*(\Lambda)$ is simple by [23, Theorem 3.1]. Thus $C^*(G_\Lambda) = C_r^*(G_\Lambda)$ is simple and hence G_Λ is topologically principal and minimal by [4, Theorem 5.1].

We have that the collection of cylinder sets of the form $Z(\mu)$ form a basis $\mathcal{B}$ of $G_\Lambda^{(0)}$ consisting of compact open sets. Now we apply Theorem 4.1 to see that $C^*(G_\Lambda)$ is purely infinite if and only if each $\chi_{Z(\mu)}$ is infinite. Let $\phi : C^*(\Lambda) \to C^*(G_\Lambda)$ be the isomorphism characterized by $s_\mu \mapsto \chi_{Z(\mu)}$. Since ϕ is an isomorphism, this gives $\chi_{Z(\mu)}$ is infinite if and only if $\phi^{-1}(\chi_{Z(\mu)}) = s_\mu s_\mu^*$ is infinite. Finally, $s_\mu s_\mu^*$ if and only if $s_{s(\mu)}$ is infinite by (1).

For (3), given an infinite path x, the collection of compact open sets of the form $Z(x(0,n))$ for $n \in \mathbb{N}^k$ form a neighbourhood base at x. Now proceed as in the proof of (2) replacing Theorem 4.1 with Corollary 4.2 and μ with $x(0,n)$. □

6. The non-simple case

Let A be a C^*-algebra. A pair of positive elements $(a_1, a_2) \in A \times A$ has the *matrix diagonalization property in* A in the sense of [15, Definition 3.3.] if for every positive matrix $\left(\begin{smallmatrix} a_1 & b_{12} \\ b_{21} & a_2 \end{smallmatrix}\right)$ with $b_{ij} \in A$ and every $\epsilon_1, \epsilon_2, \delta > 0$ there exists $d_1, d_2 \in A$ with

$$\|d_i^* a_i d_i - a_i\| < \epsilon_i \text{ and } \quad \|d_i^* b_{ij} d_j\| < \delta.$$

A subset $\mathcal{F}$ of A^+ is a *filling family* for A, in the sense of [15, Definition 3.10], if for every hereditary C^*-subalgebra H of A and every primitive ideal I of A with $H \not\subseteq I$ there exist $f \in \mathcal{F}$ and $z \in A$ with $z^* z \in H$ and $zz^* = f \notin I$.

By Proposition 3.13 and Lemma 3.12 of [15], if A^+ contains a filling family $\mathcal{F}$ that is closed under ϵ-cut-downs and every pair of elements $(a_1, a_2) \in \mathcal{F} \times \mathcal{F}$ has the matrix diagonalization property, then A is strongly purely infinite. In this section we provide a characterization of when the reduced groupoid C^*-algebra is strongly purely infinite (Proposition 6.3). In our proof of Proposition 6.3 we will use results from [4] to describe ideals of reduced groupoid C^*-algebras. First we need the following lemma. Recall that a subset $D \subseteq G^{(0)}$ is said to be *invariant* if $G \cdot D := \{r(\gamma) : s(\gamma) \in D\} \subseteq D$.

Lemma 6.1. *Let* G *be a second countable, locally compact, Hausdorff and étale groupoid such that* $C^*(G) = C_r^*(G)$. *Then the following properties are equivalent:*

(1) *For every closed invariant set* $D \subseteq G^{(0)}$

$$C^*(G|_D) = C_r^*(G|_D).$$

(2) *For every closed invariant set* $D \subseteq G^{(0)}$ *the sequence*

$$0 \longrightarrow C_r^*(G|_{G^{(0)}-D}) \xrightarrow{\iota_r} C_r^*(G) \xrightarrow{\rho_r} C_r^*(G|_D) \longrightarrow 0$$

is exact where ι_r *and* ρ_r *are determined on continuous functions by extension and restriction respectively.*

Remark 6.2. In [22, Remark 4.10], Renault mentions that if $G|_D$ is amenable for every closed invariant set $D \subseteq G^{(0)}$, then item (2) of Lemma 6.1 follows. Thus Lemma 6.1 is a strengthening of Renault's comment.

Proof. Fix a closed invariant set $D \subseteq G^{(0)}$ and let $U = G^{(0)} - D$. Consider the following diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C^*(G|_U) & \xrightarrow{\iota} & C^*(G) & \xrightarrow{\rho} & C^*(G|_D) & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle \pi_U} & & \downarrow{\scriptstyle \pi} & & \downarrow{\scriptstyle \pi_D} & & \\ 0 & \longrightarrow & C_r^*(G|_U) & \xrightarrow{\iota_r} & C_r^*(G) & \xrightarrow{\rho_r} & C_r^*(G|_D) & \longrightarrow & 0 \end{array} \tag{6.1}$$

where π_U, π and π_D are the respective quotient maps, and ι, ι_r and ρ, ρ_r extend extension and restriction respectively. Since all of the maps involved are continuous, the diagram commutes. We also have that the top row of (6.1) is exact by [18, Lemma 2.10].

(2)⇒(1): We show the surjective map π_D is injective. Fix any $a \in C_r^*(G|_D)$ with $\pi_D(a) = 0$. Find $b \in C^*(G)$ with $\rho(b) = a$. From $\pi_D(\rho(b)) = \rho_r(\pi(b)) = 0$, exactness of (6.1), surjectivity of π_U, and $\iota \circ \pi_U = \pi \circ \iota$ we obtain

$$\pi(b) \in \ker \rho_r = \iota_r(C_r^*(G|_U)) = \iota_r \circ \pi_U(C^*(G|_U)) = \pi \circ \iota(C^*(G|_U)).$$

Find $c \in C^*(G|_U)$ with $\pi(b) = \pi \circ \iota(c)$. As π is an isomorphism by assumption we obtain that $b = \iota(c)$. Hence $a = \rho(b) = \rho \circ \iota(c) = 0$, and $C^*(G|_D) = C_r^*(G|_D)$.

(1)⇒(2): By assumption the maps π and π_D are isomorphisms. Using the commutative diagram (6.1) and the exactness of the top line of that diagram, the exactness of the bottom line follows from a easy diagram chase. □

Let G be a second countable, locally compact, Hausdorff and étale groupoid and D be a closed invariant set of $G^{(0)}$ and define $U = G^{(0)} - D$. Recall from Lemma 2.1(4) we have the communing diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C_r^*(G|_U) & \xrightarrow{\iota_r} & C_r^*(G) & \xrightarrow{\rho_r} & C_r^*(G|_D) & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle E_U} & & \downarrow{\scriptstyle E} & & \downarrow{\scriptstyle E_D} & & \\ 0 & \longrightarrow & C_0(U) & \xrightarrow{\iota_0} & C_0(G^{(0)}) & \xrightarrow{\rho_0} & C_0(D) & \longrightarrow & 0. \end{array} \tag{6.2}$$

Notice that the bottom row in (6.2) is exact. We will use this diagram several times. We also use the notation Ideal$[S]$ for the ideal in $C_r^*(G)$ generated by $S \subseteq C_r^*(G)$.

Proposition 6.3. *Let G be a second countable, locally compact, Hausdorff and étale groupoid such that $C^*(G) = C_r^*(G)$. Then the following properties are equivalent:*

(1) *The C^*-algebra $C_r^*(G)$ is strongly purely infinite, and for every ideal I in $C_r^*(G)$,*

$$I = \text{Ideal}[I \cap C_0(G^{(0)})].$$

(2) *For every closed invariant set $D \subseteq G^{(0)}$, $G|_D$ is topologically principal; the sequence*

$$0 \longrightarrow C_r^*(G|_U) \xrightarrow{\iota_r} C_r^*(G) \xrightarrow{\rho_r} C_r^*(G|_D) \longrightarrow 0 \tag{6.3}$$

is exact where $U = G^{(0)} - D$, ι_r and ρ_r are determined on continuous functions by extension and restriction respectively; and for every pair of elements a, b in $C_0(G^{(0)})^+$ the 2-tuple (a, b) has the matrix diagonalization property in $C_r^*(G)$.

Proof. (1)⇒(2): Fix a closed invariant set $D \subseteq G^{(0)}$ and $U = G^{(0)} - D$. For this D and U we have a commuting diagram as in (6.2). Define $I := \ker \rho_r \subseteq C_r^*(G)$. Using the diagram, $\rho_0(E(I)) = E_D(\rho_r(I)) = 0$, implying that $E(I) \subseteq \iota_0(C_0(U))$. Since $E(b) = b$ for $b \in C_0(G^{(0)})$, $I \cap C_0(G^{(0)}) \subseteq E(I)$. Using assumption (1) we have $I = \text{Ideal}[I \cap C_0(G^{(0)})]$. Hence

$$\ker \rho_r = I = \text{Ideal}[I \cap C_0(G^{(0)})] \subseteq \text{Ideal}[E(I)] \subseteq \text{Ideal}[\iota_0(C_0(U))] \subseteq \iota_r(C_r^*(G|_U));$$

that is $\ker \rho_r \subseteq \text{image}(\iota_r)$. Thus (6.3) is exact.

We know that each $G|_D$ is topologically principal by [4, Remark 5.10] provided that $C^*(G|_D) = C_r^*(G|_D)$. The latter follows from Lemma 6.1 since (6.3) is exact.

Since $C_r^*(G)$ is strongly purely infinite, Lemma 5.8 in [13] implies that every pair (a, b) of positive elements in $C_0(G^{(0)})$ has the matrix diagonalization property in $C_r^*(G)$.

(2)⇒(1): Since we assumed that $G|_D$ is topologically principal for all closed invariant $D \subseteq G^{(0)}$, by the proof of Corollary 5.9 in [4] (see also [4, Remark 5.10]), we know $I = \text{Ideal}[I \cap C_0(G^{(0)})]$ for every ideal I in $C_r^*(G) = C^*(G)$ provided that $C^*(G|_D) = C_r^*(G|_D)$ for every closed invariant set $D \subseteq G^{(0)}$. But this follows from Lemma 6.1 since $C^*(G) = C_r^*(G)$ and (6.3) is exact, which are assumed in (2). Hence (2) implies $I = \text{Ideal}[I \cap C_0(G^{(0)})]$.

We prove $C_r^*(G)$ is strongly purely infinite. Define $\mathcal{F} := C_0(G^{(0)})^+ \subseteq C_r^*(G)$. By functional calculus we know

$$f(a) \in \mathcal{F}, \quad \text{for} \quad f \in C_0(\mathbb{R})^+, \quad a \in \mathcal{F}.$$

In particular $\mathcal{F}$ is closed under ε-cut-downs, i.e., for each $a \in \mathcal{F}$, and $\varepsilon \in (0, \|a\|)$ we have $(a - \varepsilon)_+ \in \mathcal{F}$. By (2) each pair (a, b) with $a, b \in \mathcal{F}$ has the matrix diagonalization property (of [15, Definition 3.3]). Now by Lemma 3.12 of [15] we know that $\mathcal{F}$ has the matrix diagonalization property of [15, Definition 3.10(ii)]. If follows from Proposition 3.13 of [15] that $C_r^*(G)$ is strongly purely infinite provided that $\mathcal{F}$ is a filling family for $C_r^*(G)$, which we now show.

Fix any hereditary C^*-subalgebra H of $C_r^*(G)$ and any ideal I of $C_r^*(G)$ with $H \not\subseteq I$. We know $I = \text{Ideal}[I \cap C_0(G^{(0)})]$, hence $I = \iota_r(C_r^*(G|_U))$ for some open invariant set $U \subseteq G^{(0)}$. Let $D = G^{(0)} - U$ and consider the corresponding commuting diagram (6.2).

Select $d \in H^+$, $d \notin I$. Define $c := \rho_r(d)$. As $d \notin I = \ker \rho_r$ by exactness in (2), we know $\rho(d) \neq 0$. Since E_D is faithful and c positive,

$$\epsilon := \frac{1}{4}\|E_D(c)\| > 0.$$

By (2) the groupoid $G|_D$ is topologically principal, hence Lemma 3.1 gives $f \in C_0(D)^+$ such that

$$\|f\| = 1, \quad \|fcf - fE_D(c)f\| < \epsilon, \quad \|fE_D(c)f\| > \|E_D(c)\| - \epsilon.$$

Recall [13, Lemma 2.2]: For x, y positive and $\delta > 0$ with $\|x - y\| < \delta$ there exist a contraction a with $a^*xa = (y - \delta)_+$. Use this to find a contraction $a \in C_r^*(G|_D)$ such that

$$h := a^*fcfa = (fE_D(c)f - \epsilon)_+ \in C_0(D)^+.$$

Notice that

$$\|h\| \geq \|fE_D(c)f\| - \epsilon > \|E_D(c)\| - 2\epsilon > 0.$$

Using that ρ_r restricts to the map $C_0(G^{(0)}) \to C_0(D)$, select $b \in C_0(G^{(0)})^+$ such that $\rho_r(b) = h$. Also as ρ_r is surjective find $w \in C_r^*(G)$ such that $\rho_r(w) = fa$. Since $\rho_r(b -$

$w^*dw) = h - a^*fcfa = 0$ we have $b = w^*dw + v$ for some $v \in I$. Let $\{e_\lambda\}$ denote an approximate unit of $I = C_r^*(G|_U)$ with $e_\lambda \in C_0(U)$ (see Lemma 2.1). Let 1 be the unit of the unitisation of $C_r^*(G)$. Then $(1 - e_\lambda)v(1 - e_\lambda) \to 0$. For suitable λ_0 and $e := 1 - e_{\lambda_0}$ we get

$$\|ew^*dwe - ebe\| = \|eve\| < \epsilon.$$

Use Lemma 2.2 of [13] to find a contraction $u \in C_r^*(G)$ such that

$$g := u^*ew^*dweu = (ebe - \epsilon)_+ \in C_0(G^{(0)})^+ = \mathcal{F}.$$

Since $be_{\lambda_0} + e_{\lambda_0}b - e_{\lambda_0}be_{\lambda_0} \in C_0(U) \subseteq \ker \rho_r$ we obtain that $\rho_r(ebe) = \rho_r(b) = h$. Moreover by functional calculus we know $(h - \epsilon)_+ = (fE_D(c)f - 2\epsilon)_+$. We conclude

$$\|\rho_r(g)\| = \|(h - \epsilon)_+\| = \|(fE_D(c)f - 2\epsilon)_+\| \geq \|fE_D(c)f\| - 2\epsilon > \|E_D(c)\| - 3\epsilon > 0,$$

ensuring $g \notin I$. Finally with $z := u^*ew^*d^{1/2} \in C_r^*(G)$ we obtain $g = zz^*$ and $z^*z \in H$. By definition $\mathcal{F}$ is a filling family for $C_r^*(G)$ completing the proof. □

References

[1] C. Anantharaman-Delaroche, *Purely infinite C^*-algebras arising from dynamical system*, Bull. Soc. Math. France **125** (1997), no. 2, 199-225.

[2] C. Anantharaman-Delaroche and J. Renault, *Amenable groupoids*, Monographies de L'Enseignement Mathématique, 36, L'Enseignement Mathématique, Geneva, 2000.

[3] B. Blackadar, *Operator algebras*, Encyclopaedia of Mathematical Sciences, vol. 122, Springer-Verlag, Berlin, 2006, Theory of C^*-algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III.

[4] J.H. Brown, L.O. Clark, C. Farthing and A. Sims, *Simplicity of algebras associated to étale groupoids*, Semigroup forum, to appear.

[5] T. Carlsen and K. Thomsen, *The structure of the C^*-algebra of a locally injective surjection*, Ergodic Theory Dynam. Systems **32** (2012), no. 4, 1226–1248.

[6] L.O. Clark, C. Farthing, A. Sims and M. Tomforde, *A groupoid generalization of Leavitt path algebras*, Semigroup forum, to appear.

[7] J. Cuntz, *K-theory for certain C^*-algebras*, Ann. of Math. (2) **113** (1981), no. 1, 181-197.

[8] G. Elliott, *On the classification of C^*-algebras of real rank zero*, J. Reine Angew. Math. **443** (1993), 179-219.

[9] R. Exel, *Reconstructing a totally disconnected groupoid from its ample semigroup*, Proc. Amer. Math. Soc. **138** (2010), 2991–3001.

[10] T. Giordano and A. Sierakowski, *Purely infinite partial crossed products*, submitted. arXiv:1303.4483v2

[11] E. Kirchberg, *The classification of purely infinite C^*-algebras using Kasparov's theory*, preprint, 1994.

[12] E. Kirchberg, *Das nicht-kommutative Michael-Auswahlprinzip und die Klassifikation nicht-einfacher Algebren*, p. 92–141 in *C*–Algebras*: Proceedings of the SFB-Workshop on C*-algebras, Münster, Germany, March 8-12, 1999/J. Cuntz, S. Echterhoff (ed.), Springer, Berlin etc. (2000), 272 pp.

[13] E. Kirchberg and M. Rørdam, *Infinite non-simple C*-algebras: absorbing the Cuntz algebras $\mathcal{O}_\infty$*, Adv. Math. **167** (2002), 195–264.

[14] E. Kirchberg and M. Rørdam, *Non-simple purely infinite C^*-algebras*, Amer. J. Math. **122** (2000), 637–666.

[15] E. Kirchberg and A. Sierakowski, *Strong pure infiniteness of crossed products,* J. Funct. Anal. (2014).

[16] A. Kumjian and D. Pask, *Higher rank graph C^*-algebras*, New York J. Math. **6** (2000), 1–20.

[17] A. Kumjian, D. Pask, and I. Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. **184** (1998), no. 1, 161–174.

[18] P.S. Muhly, J. Renault and D.P. Williams, *Continuous trace groupoid C^*-algebras, III*, Trans. Amer. Math. Soc. **348** (1996), 3621–3641.

[19] A. Paterson, *Groupoids, Inverse Semigroups, and their Operator Algebras*, Progress in Mathematics, 170, Birkhauser, Boston, 1998.
[20] N. C. Phillips, *A classification theorem for nuclear purely infinite simple C^*-algebras*, Doc. Math. **5** (2000), 49-114.
[21] J. Renault, *A groupoid approach to C^*-algebras.* Lecture Notes in Mathematics, 793. Springer, Berlin, 1980.
[22] J. Renault, *The ideal structure of groupoid crossed product C^*-algebras*, J. Operator Theory **25** (1991), 3–36.
[23] D.I. Robertson and A. Sims, *Simplicity of C^*-algebras associated to higher-rank graphs*, Bull. Lond. Math. Soc. **39** (2007), 337–344.
[24] M. Rørdam and A. Sierakowski, *Purely infinite C^*-algebras arising from crossed products*, Ergodic Theory Dynam. Systems **32** (2012), no. 1, 273-293.
[25] A. Sims, *Gauge-invariant ideals in the C^*-algebras of finitely aligned higher-rank graphs*, Canad. J. Math. **58** (2006), 1268-1290.
[26] J. Spielberg, *Graph-based models for Kirchberg algebras*, J. Operator Theory **57** (2007), 347–374.
[27] B. Steinberg, *A groupoid approach to inverse semigroup algebras*, Adv. Math. **223** (2010), 689-727.

Jonathan Brown, Mathematics Department, Kansas State University, 138 Cardwell Hall, Manhattan, KS 66506-2602, USA.
E-mail address: brownjh@math.kansas.edu

Lisa Orloff Clark, Department of Mathematics and Statistics, University of Otago, PO Box 56, Dunedin Dunedin 9054, New Zealand.
E-mail address: lclark@maths.otago.ac.nz

Adam Sierakowski, School of Mathematics and Applied Statistics, University of Wollongong, NSW 2522, Australia
E-mail address: asierakowski@uow.edu.au